

AD-754 306

ENDOGENOUS ENDOTOXEMIA DURING HEMOR-
RHAGIC SHOCK IN THE SUBHUMAN PRIMATE

Clifford M. Herman, et al

Naval Medical Research Institute
Bethesda, Maryland


1972

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151

DISCLAIMER NOTICE



**THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE COPY
FURNISHED TO DTIC CONTAINED
A SIGNIFICANT NUMBER OF
PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**


UNCLASSIFIED

AD-754 306

DOCUMENT CONTROL DATA - R & D

1. ORIGINATING AGENCY (Corporate author)			2a. REPORT SECURITY CLASSIFICATION	
NAVAL MEDICAL RESEARCH INSTITUTE NATIONAL NAVAL MEDICAL CENTER BETHESDA, MARYLAND 20014			UNCLASSIFIED	
3. REPORT TITLE			2b. GROUP	
ENDOGENOUS ENDOTOXEMIA DURING HEMORRHAGIC SHOCK IN THE SUBHUMAN PRIMATE				
4. DESCRIPTIVE NOTES (Type of report and, inclusive dates)				
MEDICAL RESEARCH PROGRESS REPORT				
5. AUTHOR(S) (First name, middle initial, last name)				
Clifford M. Herman, Avram R. Kraft, Kenneth R. Smith, E. Joseph Artnak, Fleming C. Chisholm, Larry G. Dickson and Louis D. Homer				
6. REPORT DATE		7a. TOTAL NO. OF PAGES		7b. NO. OF REFS
1972		2		1
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)		
b. PROJECT NO.		M4305.05.3007BGGO Report No. Report No. 22		
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)		
d.				
10. DISTRIBUTION STATEMENT				
THIS DOCUMENT HAS BEEN APPROVED FOR PUBLIC RELEASE AND SALE: ITS DISTRIBUTION IS UNLIMITED				
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY		
Reprinted from: Surgical Forum Volume XXIII, 1972		Bureau of Medicine and Surgery (Navy) Washington, D. C. 20372		
13. ABSTRACT				
Ischemic bowel damage resulting in endogenous endotoxemia is not a common feature of hemorrhagic shock in baboons. The observed sporadic endotoxemia occurs no more frequently during shock than in the baseline period, and is not related to the duration or degree of severity of hemorrhagic shock in this subhuman primate species.				

NATIONAL TECHNICAL
INFORMATION SERVICE
Department of Commerce
Springfield, MA 01104



ENDOGENOUS ENDOTOXEMIA DURING HEMORRHAGIC SHOCK IN THE SUBHUMAN PRIMATE

CLIFFORD M. HERMAN, MD, FACS, AVRAM R. KRAFT, MD,
KENNETH R. SMITH, BS, E. JOSEPH ARTNAK, BS,
FLEMING C. CHISHOLM, BS, LARRY G. DICKSON, MD, AND
LOUIS D. HOMER, MD, PhD

AD754306

EXPERIMENTS were carried out to test the hypothesis that, during hemorrhagic shock, endotoxin enters the portal circulation from ischemic bowel and is then associated with irreversibility of the hemorrhagic shock state. Since this theory has evolved mainly from canine experimental data, we used baboons in order to determine the validity of the theory in a species phylogenetically closer to man.

METHODS

After placement of sampling catheters in the portal vein, right atrium, and aorta, 14 awake, restrained baboons were subjected to 1 hr of hemorrhagic shock at a mean arterial pressure (MAP) of 60 torr followed by a second hour at 40-torr MAP. Six animals were resuscitated with lactated Ringer's solution in a volume three times the total blood loss followed by reinfusion of their shed blood. Eight animals were maintained hypotensive until death. Serial blood samples from all three sites were analyzed for endotoxin by the Limulus lysate method (1). Comparisons of incidence of endotoxemia according to sampling site and period of shock were made by Chi-square analysis.

RESULTS

The assay was 100% effective in detecting endotoxin at a concentration of 10^{-2} mg/ml, 94% effective at 10^{-4} mg/ml, and 39% effective at 10^{-6} mg/ml. There were no false-positive results.

From the Division of Experimental Surgery, Naval Medical Research Institute, National Naval Medical Center, Bethesda, Md. Supported by Navy Department Research Task M4305.05.3007BGGO.

Reprint from Surgical Forum Volume XXIII, 1972

Table 1

	BASLINE*	EARLY SHOCK* (MAP > 40 TORR)	LATE SHOCK* (MAP < 40 TORR)
Resuscitated group	5/36	6/36	6/36
Non-resuscitated group	5/48	3/48	6/108

* Values expressed as samples positive for endotoxin / total number of samples.

Endotoxemia was found infrequently (Table 1), with no greater incidence ($P > .6$) in portal venous samples than in systemic blood, so these data were pooled for further analysis. Furthermore, endotoxemia was no more frequent ($P > .6$) late in shock than it was in early shock or during the baseline period. Autopsy showed only mild submucosal edema of the bowel in the resuscitated animals, with no splanchnic or other organ abnormalities in the nonresuscitated group.

COMMENT

Several possible explanations can be considered for the inconsistent and infrequent finding of endotoxemia in this study. Endotoxin could have been present at concentrations below the limit of sensitivity of the detection method. However, while it is not known what level of endotoxin can be damaging during a low flow state, endotoxin concentrations which are lethal in themselves in a baboon would have been detected with 100% certainty by our assay method. With no signs of ischemic bowel damage, it is more likely that the mesenteric blood flow and the integrity of the splanchnic viscera are not selectively compromised by hemorrhagic shock in this subhuman primate species.

CONCLUSION

Ischemic bowel damage resulting in endogenous endotoxemia is not a common feature of hemorrhagic shock in baboons. The observed sporadic endotoxemia occurs no more frequently during shock than in the baseline period, and is not related to the duration or degree of severity of hemorrhagic shock in this subhuman primate species.

REFERENCE

1. Levin J, Tomasulo PA, Oser RS: Detection of endotoxin in human blood and demonstration of an inhibitor. *J Lab Clin Med* 75:903-908, 1970